arXiv:1405.7757v1 [math.OA] 30 May 2014

AF-Embeddings of Graph Algebras

Christopher Schafhauser

Abstract

Let E be a countable directed graph. We show that $C^*(E)$ is AF-embeddable if and only if no loop in E has an entrance. The proof is constructive and is in the same spirit as the Drinen-Tomforde desingularization in [4].

Introduction

In [7], Pimnser and Voiculescu argued the irrational rotation algebras A_θ can be embedded into an AF C^*-algebra. Since then, there has been an interest in characterizing the C^*-algebras which are AF-embeddable; especially crossed products. Pimnser [6] and Brown [2], repsectively, have solved the AF-embeddability question for algebras of the form $C(X) \rtimes \mathbb{Z}$ for a compact metric space X and $A \rtimes \mathbb{Z}$ for an AF-algebra A. See [3, Chapter 8] for a survey on AF-embeddability.

The general AF-embeddability problem is still largely unsolved. There are only two known obstructions to AF-embeddability; namely exactness and quasidiagonality. A C^*-algebra A is said to be *exact*, if the functor $B \mapsto A \otimes_{\min} B$ preserves short exact sequences. A C^*-algebra is called *quasidiagonal* if there are sequences of finite dimensional C^*-algebras F_n and completely positive contractive maps $\varphi_n : A \to F_n$ such that

$$\|\varphi_n(ab) - \varphi_n(a)\varphi_n(b)\| \to 0 \qquad \text{and} \qquad \|\varphi_n(a)\| \to \|a\|$$

for every $a, b \in A$. See [3, Chapters 3 and 7] for an introduction to exactness and quasidiagonality.

Both quasidiagonality and exactness are preserved by taking subalgebras and AF-algebras enjoy both properties. Hence every AF-embeddable C^*-algebra is exact and quasidiagonal. It is conjectured

in [1] that the converse is true. Blackadar and Kirchberg also ask if every stably finite nuclear C^*-algebra is quasidiagonal. Hence in particular, the conjecture is that stable finiteness, quasidiagonality, and AF-embeddability are equivalent for nuclear C^*-algebras. The main result of this paper verifies this conjecture for graph C^*-algebras. In particular, we have

Theorem 1. *For a countable graph E, the following are equivalent:*

1. *$C^*(E)$ is AF-embeddable;*
2. *$C^*(E)$ is quasidiagonal;*
3. *$C^*(E)$ is stably finite;*
4. *$C^*(E)$ is finite;*
5. *No loop in E has an entrance.*

Graph C^*-Algebras

By a graph we mean a quadruple $E = (E^0, E^1, r, s)$, where E^0 and E^1 are countable sets called the *vertices* and *edges* of E, and $r, s : E^1 \to E^0$ are functions called the *range* and *source* maps. Given a graph E, a Cuntz-Krieger E-family in a C^*-algebra A is a collection

$$\{p_v, s_e : v \in E^0, e \in E^1\} \subseteq A$$

such that for all $v \in E^0$ and $e, f \in E^1$, we have

1. $p_v^2 = p_v = p_v^*$ for all $v \in E^0$
2. $s_e^* s_f = \begin{cases} p_{s(e)} & e = f \\ 0 & e \neq f \end{cases}$
3. $p_v = \sum_{e \in r^{-1}(v)} s_e s_e^*$ if $0 < |r^{-1}(v)| < \infty$.

Let $C^*(E)$ denote the universal C^*-algebra generated by a Cuntz-Krieger E-family. See [8] for an introduction to graph C^*-algebras.

If E is a graph and $n \geq 1$, a path in E is a list of edges $\alpha = (\alpha_n, \ldots, \alpha_1)$ such that $r(\alpha_i) = s(\alpha_{i+1})$ for each $1 \leq i < n$. Define $r(\alpha) = r(\alpha_n)$ and $s(\alpha) = s(\alpha_1)$. Define E^n to be the set of paths of length n in E and $E^* = \bigcup_{n=0}^{\infty} E^n$ the paths of finite length in E. In

particular, the vertices of E are considered to be paths of length 0. Given $\alpha = (\alpha_n, \ldots, \alpha_1)$, define $s_\alpha = s_{\alpha_n} \cdots s_{\alpha_1}$. It can be shown that

$$C^*(E) = \overline{\text{span}}\{s_\alpha s_\beta^* : \alpha, \beta \in E^* \text{ with } s(\alpha) = s(\beta)\}.$$

A loop in E is a path $\alpha \in E^n$ with $n \geq 1$ such that $r(\alpha) = s(\alpha)$. We say α is a *simple loop* if $r(\alpha_i) \neq r(\alpha_j)$ for $i \neq j$. We say α has an entrance if $|r^{-1}(r(\alpha_i))| > 1$ for some i. The structure of the algebra $C^*(E)$ is closely related to the structure of the loops in E. We will show in Theorem 1, the AF-embeddability of $C^*(E)$ is also characterized by the loops in E.

We recall two results about graph C^*-algebras. Theorem 2 is from Kumijan, Pask, and Raeburn in the row-finite case and Drinen and Tomforde in general (see [5, Theorem 2.4] and [4, Corollary 2.13]). Theorem 3 is Szymański's generalization of the Cuntz-Krieger Uniqueness Theorem (see [9, Theorem 1.2]).

Theorem 2. *For a countable graph E, $C^*(E)$ is AF if and only if E has no loops.*

Theorem 3. *Suppose E is a graph, A is a C^*-algebra, and $\{\tilde{p}_v, \tilde{s}_e\} \subseteq A$ is a Cuntz-Kreiger E-family. If $\tilde{p}_v \neq 0$ for every $v \in E^0$ and $\sigma(\tilde{s}_\alpha) \supseteq \mathbb{T}$ for every entry-less loop $\alpha \in E^*$, then the induced morphism $C^*(E) \to A$ defined by $p_v \mapsto \tilde{p}_v$ and $s_e \mapsto \tilde{s}_e$ is injective.*

Proof of Theorem 1

We are now ready to prove our main result. Starting with a graph E satisfying condition (5), we will replace each loop in E with the Bratteli diagram of an AF-algebra to build a new graph F such that $C^*(F)$ is AF and $C^*(E) \subseteq C^*(F)$. The idea of the proof is motivated by the Drinen-Tomforde desingularization process introduced in [4].

Proof of Theorem 1. It is well-known that (1) implies (2) and (2) implies (3) (see [3, Propositions 7.1.9, 7.1.10, and 7.1.15]) and it is obvious that (3) implies (4). To see (4) implies (5), note that if $\alpha, \beta \in E^*$ are distinct paths with $s(\alpha) = r(\alpha) = r(\beta)$, then we have

$$s_\alpha^* s_\alpha = p_{s(\alpha)} \quad \text{and} \quad s_\alpha s_\alpha^* \lneq s_\alpha s_\alpha^* + s_\beta s_\beta^* \leq p_{s(\alpha)}.$$

So $p_{s(\alpha)}$ is an infinite projection and $C^*(E)$ is infinite.

Now suppose (5) holds. Choose a unital AF-algebra A such that there is a unitary $t \in A$ with $\sigma(t) = \mathbb{T}$ and let B be a Bratteli diagram for A with sink v. Let $e_n \cdots e_2 e_1$ be a simple loop in E and set $u_i = s(e_i)$. Define a graph F by

$$F^0 = E^0 \cup B^0, \qquad F^1 = \big(E^1 \setminus \{e_1, \ldots, e_n\}\big) \cup B^1 \cup \{f_1, \ldots f_n\}$$

and extend the range and source maps by $r(f_i) = u_i$ and $s(f_i) = v$. For example, if $A = M_{2^\infty}$, and E and B the graphs



then F is the graph given below:



Note that $p_v C^*(F) p_v \cong A$ and hence we may view t as an element of $C^*(F)$. Define $\tilde{s}_{e_i} = s_{f_{i+1}} t s_{f_i}^* \in C^*(F)$ for each $i = 1, \ldots, n$. Since no loop in E has an entrance, we have $r_F^{-1}(u_i) = \{f_i\}$. Hence

$$\tilde{s}_{e_i}^* \tilde{s}_{e_i} = s_{f_i} s_{f_i}^* = p_{u_i} \quad \text{and} \quad \tilde{s}_{e_i} \tilde{s}_{e_i}^* = s_{f_{i+1}} s_{f_{i+1}}^* = p_{u_{i+1}}.$$

Moreover,

$$\sigma(\tilde{s}_{e_n} \tilde{s}_{e_{n-1}} \cdots \tilde{s}_{e_1}) = \sigma(s_{f_1} t^n s_{f_1}^*) = \sigma(s_{f_1}^* s_{f_1} t^n) = \sigma(t^n) = \mathbb{T} \cup \{0\}.$$

Now, by Theorem 3, there is an inclusion $C^*(E) \hookrightarrow C^*(F)$ given by

$$p_v \mapsto p_v \text{ for } v \in E^0 \quad \text{and} \quad s_e \mapsto \begin{cases} \tilde{s}_e & e \in \{e_1, \ldots, e_n\}, \\ s_e & e \in E^1 \setminus \{e_1, \ldots, e_n\}. \end{cases}$$

Note that since no loop in E has an entrance, the loops in the graph E are disjoint. Thus by applying the construction above to every loop in E, we may build a graph F with no loops and an embedding $C^*(E) \hookrightarrow C^*(F)$. Since F has no loops, $C^*(F)$ is AF by Theorem 2 and hence $C^*(E)$ is AF-embeddable. □

References

[1] B. Blackadar and E. Kirchberg, *Generalized inductive limits of nitedimensional C*-algebras*, Math. Ann. 307 (1997), 343 - 380.

[2] N. P. Brown, *AF Embeddability of Crossed Products of AF Algebras by the Integers*, J. Funct. Anal. **160** (1998), 150-175.

[3] N. P. Brown and N. Ozawa, *C*-Algebras and Finite-Dimensional Approximations*. Graduate Studies in Mathematics, 88. Amer. Math. Soc., Providence, RI, 2008.

[4] D. Drinen and M. Tomforde, *The C*-algebras of arbitrary graphs*, Rocky Mountain J. Math. 35 (2005), no. 1, 105135.

[5] A. Kumijan, D. Pask, and I. Raeburn, *Cuntz-Krieger algebras of directed graphs*, Pacific J. Math. **184** (1998), 161-174.

[6] *Embedding some transformation group C*-algebras into AF-algebras*, Ergodic Theory Dynam. Systems 3 (1983), no. 4, 613626.

[7] M. Pimsner and D. Voiculescu, *Imbedding the irrational rotation C*-algebra into an AF-algebra*, J. Operator Theory **4** (1980), no. 2, 201–210.

[8] I. Raeburn, *Graph algebras*, CBMS Regional Conference Series in Mathematics **103** Published for the Conference Board of the Mathematical Sciences, Washington D.C. by the AMS, Providence, RI (2005).

[9] W. Szymański, *General Cuntz-Krieger uniqueness theorem*, International J. Math. 13 (2002).

Dept. of Mathematics, University of Nebraska-Lincoln, Lincoln, NE, 68588-0130

E-mail address: cschafhauser2@math.unl.edu